# ES COMPETITOR SOYBEAN LATE GROUP (beginning of I) THE N°1 IN PROTEIN IN GROUP I AND **VERY EARLY** !

Hi-Pro by Lidea

# BENEFITS

#### VERY HIGH PROTEIN

• The richest variety in protein with +1,5 points/Isidor

#### VERY EARLY

• Allows early harvest before corn

#### FOOD CRITERIA

# **DENTITY CARD**

- Registered in: HU
- **Official standard in:** HU
- **Ø Breeder code:** ESG1943
- **Type of Growth:** Indeterminated
- **W** Hair color: tawny
- **V** Flower color: white
- **Uzeave shape:** pointed ovate
- **Pod color:** dark brown
- **Hilum color:** dark brown

# **CULTURE TIPS**

#### • Planting density should integrate all possible losses:

| Water<br>conditions    | Emerged<br>plants /ha | Planting<br>density /ha |
|------------------------|-----------------------|-------------------------|
| No irrigation<br>/ Dry | 450 000               | 500-560 000             |
| Irrigated<br>/ Rainy   | 350 000               | 390-440 000             |

- Distance between rows: 25 to 60 cm
- Planting depth: 2 to 4 cm
- Plant in warm soils (>10°C) with warm temperatures during the days after planting



#### www.lidea-seeds.com

### **HERBICIDES TOLERANCES**

| Pendimethaline |          |        |      |           |  |  |
|----------------|----------|--------|------|-----------|--|--|
| very low       | low      | medium | high | very high |  |  |
| Metobromuron   |          |        |      |           |  |  |
| very low       | low      | medium | high | very high |  |  |
| Metribuzine    | <u>)</u> |        | -    |           |  |  |
| very low       | low      | medium | high | very high |  |  |



The information provided in this document is for informational purposes only, and may vary according to agricultural and climate conditions, as well as cultivation techniques. Disease tolerance information applies to diseases or strains currently known in France. March 2021. Source: R&D Lidea.

#### TKW (Thousand Kernel Weight): 190 g very low low

very low

verv low

QUALITY

Protein content: 44% of Dry Matter

low

V

low

Oil content: 18,2% of Dry Matter

# AGRONOMIC CHARACTERISTICS

medium

medium

medium

high

high

high

very high

very high

very high

| Early vigor: 6           | 5,7 | _      |      |           |  |
|--------------------------|-----|--------|------|-----------|--|
|                          |     |        |      |           |  |
| very low                 | low | medium | high | very high |  |
| 1st pod height: 16,3 cm  |     |        |      |           |  |
| very low                 | low | medium | high | very high |  |
| Plant height: 111 cm     |     |        |      |           |  |
| very low                 | low | medium | high | very high |  |
| Lodging tolerance: 2,4   |     |        |      |           |  |
| very low                 | low | medium | high | very high |  |
| Pod shattering tolerance |     |        |      |           |  |
| very low                 | low | medium | high | very high |  |

### **DISEASES TOLERANCES**

| Sclerotinia |     |        |      |           |
|-------------|-----|--------|------|-----------|
| very low    | low | medium | high | very high |
| Diaporthe   |     |        |      |           |
| very low    | low | medium | high | very high |